

# 2009 H1N1 INFLUENZA VACCINE

## INACTIVATED (the "flu shot")

### WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis).

#### 1 What is 2009 H1N1 influenza?

2009 H1N1 influenza (also called Swine Flu) is caused by a new strain of influenza virus. It has spread to many countries.

Like other flu viruses, 2009 H1N1 spreads from person to person through coughing, sneezing, and sometimes through touching objects contaminated with the virus.

Signs of 2009 H1N1 can include:

- Fatigue
- Fever
- Sore Throat
- Muscle Aches
- Chills
- Coughing
- Sneezing

Some people also have diarrhea and vomiting.

Most people feel better within a week. But some people get pneumonia or other serious illnesses. Some people have to be hospitalized and some die.

#### 2 How is 2009 H1N1 different from regular (seasonal) flu?

Seasonal flu viruses change from year to year, but they are closely related to each other.

People who have had flu infections in the past usually have some immunity to seasonal flu viruses (their bodies have built up some ability to fight off the viruses).

The 2009 H1N1 flu is a new flu virus. It is very different from seasonal flu viruses.

Most people have little or no immunity to 2009 H1N1 flu (their bodies are not prepared to fight off the virus).

#### 3 2009 H1N1 influenza vaccine

Vaccines are available to protect against 2009 H1N1 influenza.

- These vaccines are made just like seasonal flu vaccines.
- They are expected to be as safe and effective as seasonal flu vaccines.
- They will not prevent "influenza-like" illnesses caused by other viruses.
- They will not prevent seasonal flu. *You should also get seasonal influenza vaccine, if you want to be protected against seasonal flu.*

**Inactivated** vaccine (vaccine that has killed virus in it) is injected into the muscle, like the annual flu shot. **This sheet describes the inactivated vaccine.**

A **live, intranasal** vaccine (the nasal spray vaccine) is also available. It is described in a separate sheet.

Some inactivated 2009 H1N1 vaccine contains a preservative called thimerosal to keep it free from germs. Some people have suggested that thimerosal might be related to autism. In 2004 a group of experts at the Institute of Medicine reviewed many studies looking into this theory, and found no association between thimerosal and autism. Additional studies since then reached the same conclusion.

#### 4 Who should get 2009 H1N1 influenza vaccine and when?

##### WHO

Groups recommended to receive 2009 H1N1 vaccine first are:

- Pregnant women
- People who live with or care for infants younger than 6 months of age
- Health care and emergency medical personnel
- Anyone from 6 months through 24 years of age
- Anyone from 25 through 64 years of age with certain chronic medical conditions or a weakened immune system

As more vaccine becomes available, these groups should also be vaccinated:

- Healthy 25 through 64 year olds
- Adults 65 years and older

The Federal government is providing this vaccine for receipt on a voluntary basis. However, state law or employers may require vaccination for certain persons.

##### WHEN

Get vaccinated as soon as the vaccine is available.

Children through 9 years of age should get **two doses** of vaccine, about a month apart. Older children and adults need only one dose.